

RADIESSE[®] + MEN

VOLUMIZING FILLER

196 HANGOVERS

4 HEARTBREAKS


135 SUNBURNS

3 TEENAGERS


HIDE THE EVIDENCE.

Let's face it, you earned every one of those wrinkles and creases. But that doesn't mean you have to live with them. Not when there's RADIESSE[®] Volumizing Filler.

Made from an exclusive natural mineral, RADIESSE[®] Volumizing Filler is strong enough for a man's rugged skin. After it's injected under the skin, it stimulates your body's own collagen¹, resulting in a natural look and feel. Results improve over time and may last more than 12 months in many patients.^{2,3}



BEFORE



AFTER*

*Individual results may vary.

Go to RADIESSEMEN.COM to find a provider near you to hide your evidence!

Actual
RADIESSE[®]
Patient
Age 57

MERZ AESTHETICS[™]

IMPORTANT SAFETY INFORMATION FOR RADIESSE[®] VOLUMIZING FILLER

Indication: RADIESSE[®] Volumizing Filler is FDA-approved for subdermal implantation for the correction of moderate to severe facial wrinkles and folds, such as nasolabial folds.

Contraindications: RADIESSE[®] Volumizing Filler should not be used in patients with bleeding disorders or in patients with severe allergies manifested by a history of anaphylaxis, with a history or presence of multiple severe allergies or with a history of hypersensitivity to the components of RADIESSE[®] Volumizing Filler.

Warnings: RADIESSE[®] Volumizing Filler should not be injected into blood vessels. Use of RADIESSE[®] Volumizing Filler in any person with active skin inflammation or infection in or near the treatment area should be deferred until the inflammatory or infectious process has been controlled. The safety and effectiveness for use in the lips has not been established. There have been published reports of nodules associated with the use of RADIESSE[®] Volumizing Filler injected into the lips.

Precautions: Safety for use during pregnancy, in breastfeeding females or in patients under 18 years has not been established. As with all skin-injection procedures, there is a risk of infection. Patients using medications that prolong bleeding, such as aspirin or warfarin, may, as with any injection procedure, experience increased bruising and bleeding at the injection site. Patients should inform their physician if they are using such medications. Patients should minimize exposure of the treated area to extensive sun or heat exposure for approximately 24 hours after treatment or until any initial swelling and redness has resolved. Safety and effectiveness in the periorbital area has not been established. The safety of RADIESSE[®] Volumizing Filler in patients with a susceptibility to keloid formation and hypertrophic scarring has not been studied. Patients with a history of previous herpetic eruption may experience reactivation of the herpes.

Adverse Events: After injection, patients may experience redness, bruising, swelling or other local side effects. Most side effects of treatment resolve within a few days. More rare side effects may include swelling that lasts longer, unevenness or firmness in the area injected.

Important: For full safety information, please visit www.Radiesse.com or call Merz Aesthetics Customer Service at 866-862-1211

Caution: Rx Only

¹Berlin A, Hussain M, Goldberg D. (2008) Calcium Hydroxylapatite Filler (RADIESSE[®]) for Facial Rejuvenation: A Histologic and Immunohistochemical Analysis. *Dermatologic Surgery*, Volume 34, S64-S67.

²Moers-Carpi M, Vogt S, et. al. (2007) A Multicenter, Randomized Trial Comparing Calcium Hydroxylapatite to Two Hyaluronic Acids for Treatment of Nasolabial Folds. *Dermatologic Surgery*, 33 (2S), 144s-151s.

³Bass LS, Smith S, Busso M, McClaren M. (2010) Calcium Hydroxylapatite (RADIESSE[®]) for Treatment of Nasolabial Folds: Long-Term Safety and Efficacy Results. *Aesthetic Surgery J*, 30 (2): 235-238.

DERMA HEALTH

Laser & Skin Care Clinic

Confidence
You'll Notice.